

Multiple Sub-Jet Model of Gamma Ray Bursts and Possible Origin of X-ray Precursors and Post-cursors

Takashi Nakamura

Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606

ABSTRACT

We assume that internal shocks of Gamma Ray Bursts (GRBs) consist of multiple sub-jets with a collimation half-angle $\sim \text{several} \times \gamma_i^{-1}$, where γ_i is the Lorentz factor of each sub-jet. If by chance a sub-jet is first emitted off-axis from the line of sight, the observed peak energy can be in the X-ray region. Next if by chance a subsequent sub-jet is emitted along the line of sight, then the peak energy will be in the gamma ray region and the gamma ray may arrive after the X-ray precursor from the former sub-jet depending on parameters. This model predicts a new class of GRBs with extremely weak and short gamma ray emission but X-ray precursor and/or post-cursor as well as an afterglow. (ApJ Letters; Received: 2000 Jan 31 Accepted: 2000 Mar 30)

Subject headings: gamma rays: bursts

1. Introduction

Recently evidences for collimation of GRBs in the afterglow came out for GRB990123 (Kulkarni *et al.* 1999, Fruchter *et al.* 1999, Galama *et al.* 1999, Castro-Tirado *et al.* 1999; see also Dai & Lu 1999 for non-jet interpretation) and GRB990510 (Harrison *et al.* 1999, Stanek *et al.* 1999). The rapid decline rate of GRB980519 (Halpern, Kemp, Piran & Bershadsky 1999) also suggests the collimation of GRBs. From the fact that the power index of the decline rate of the afterglow changed at $1 \sim 2$ days for GRB990123 and GRB990510, the collimation half-angle $\Delta\theta_a$ of the afterglow is estimated as ~ 0.1 (Rhoads 1997, Kulkarni *et al.* 1999, Harrison *et al.* 1999), where a stands for afterglow. The unusually slowly declining X-ray afterglow of GRB980425 ($\propto T^{-0.2}$) can also be explained¹ if the afterglow is beamed and the angle between the jet and the line of sight is $\sim 30^\circ$ (Nakamura 1999).

In an internal shock model (Piran 1998), which is one of the promising models of GRBs, the relativistic beaming half-angle ($= \gamma_i^{-1} < 0.01$) with γ_i being the initial gamma factor of the internal shock is smaller than the suggested collimation half-angle of the afterglow ($\Delta\theta_a \sim 0.1$). This means that it is possible to assume that the collimation half-angle of the internal shock is much smaller than the collimation half-angle of the afterglow. We here assume that the beam of internal shocks consist of such sub-jets with the collimation half-angle $\Delta\theta_\gamma < \Delta\theta_a$. This kind of a model has already been considered to interpret no correlation between the isotropic luminosity of gamma-rays and that of the afterglow (Kumar & Piran 1999) as well as the diversity of the isotropic luminosity for GRBs with known red-shifts.

Here I like to point out that such a sub-jet model has a potential ability to interpret X-ray precursors, which were first observed by Ginga satellite (Murakami *et al.* 1991). The evidence for X-ray precursors have been confirmed by GRANAT/WATCH (Sazonov *et al.* 1998) catalogue with 6 X-ray precursor events in 95 GRBs. This suggests that about $\sim 6\%$ of GRBs have precursors. Recently Beppo/SAX found the X-ray precursor and prompt X-ray emission from GRB980519 (in't Zand *et al.* 1999). For GRB980519 an analysis of the evolution of the X-

ray spectrum in terms of a single power-law spectrum shows that the photon index evolved from -2.0 to -1.1 to -2.4. The onset of burst has such a soft spectrum that the 2-27 keV emission appears to precede the > 107 keV emission by about 70s. One more important fact for GRB980519 is that the X-ray post-cursor is seen after ~ 100 s from the gamma ray emission.

Paczynski 1998 has once suggested that precursors of GRBs might be the evidence for the association of GRBs with supernovae, that is, precursors might suggest baryon contaminated environment of GRBs, while the gamma ray might come from the clean fire ball. Here I like to interpret precursors in the framework of the standard fire ball model which is so successful so far (Piran 1998).

In this *Letter*, I assume that internal shocks of Gamma Ray Bursts (GRBs) consist of multiple sub-jets with the collimation half-angle $\sim \text{several} \times \gamma_i^{-1}$, where $\gamma_i \simeq 100 \sim 1000$ is the Lorentz factor of each sub-jet. If by chance a sub-jet is first emitted off-axis from the line of sight, the observed peak energy can be in the X-ray region. Next if by chance a subsequent sub-jet is emitted along the line of sight, then the peak energy will be in the gamma ray region and depending on parameters, the gamma ray can arrive after the X-ray precursor from the former sub-jet. This emission pattern agrees with that of GRBs with precursors. The event rate of precursors ($\sim 6\%$) is compatible with this model since precursors occur by chance. If the time order of the former and the latter sub-jets is reversed by chance, gamma-ray comes first and X-ray comes later. This emission pattern is compatible with the post-cursor. In next section we will argue a simple model in this picture.

2. Simple Model

Let us consider a simple model of GRBs in the framework of an internal shock picture. First at $t = 0$ in the laboratory frame a slow proton jet with Lorentz factor $\gamma_s > 100$ with the collimation half-angle $\Delta\theta_{\gamma_s}$ starts from the central engine toward the direction θ_v where θ_v is the angle between the direction to the detector and the axis of the sub-jet. This slow jet continues for Δt_s . At $t = \Delta t_s + t_r$, a rapid proton jet with Lorentz factor $\gamma_r > \gamma_s$ with the collimation half-angle $\Delta\theta_{\gamma_r}$ starts from the central engine toward the same direction θ_v as the slow jet. For simplicity we consider only the case of $\Delta\theta_{\gamma_s} = \Delta\theta_{\gamma_r} \equiv \Delta\theta_\gamma$. Then the rapid jet catches

¹In this Letter I use T to express the time at the detector. For simplicity I neglect the cosmological time dilatation to avoid confusions.

up the slow jet at $t_c = 2t_r\gamma_s^2 S^2/(S^2 - 1) + \Delta t_s$ where $S \equiv \gamma_r/\gamma_s > 1$. The internal shock breaks out at $t_b = t_c + 2\Delta t_s\gamma_s^2 S'^2/(S'^2 - 1)$, where $S' \equiv \gamma_{sh}/\gamma_s > 1$ with γ_{sh} being the gamma factor of the internal shock in the laboratory frame. Let us consider the case of $\theta_v > \gamma_s^{-1}$. Then the relation of the observed time T to t is given by

$$T = (1 - \beta \cos \theta_v)t \sim t\theta_v^2/2, \quad (1)$$

where we set $T = 0$ as the time when the photon starts from the central engine at $t = 0$ arrives at the detector. The observed frequency ν at the detector is related to ν' in the shock frame as

$$\nu = \frac{\nu'}{\gamma(1 - \beta \cos \theta_v)} \sim 11.1\nu'(\frac{\gamma_{sh}}{200})^{-1}(\frac{\theta_v}{0.03})^{-2}. \quad (2)$$

Since in the internal shock model the peak energy in the comoving frame should be in the soft X-ray band (say $\sim \text{keV}$), the observed peak energy from the off-axis emission can be in X-ray band (say $\sim 10\text{keV}$) for appropriate values of γ_{sh} and θ_v . For $t_r \sim \Delta t_s$, the arrival time of this off-axis emission ($T_c(\theta_v)$) is given by

$$T_c(\theta_v) = 9t_r(\frac{\gamma_s}{100})^2(\frac{\theta_v}{0.03})^2\frac{S^2}{S^2 - 1}, \quad (3)$$

while the duration from the catch up time to the shock break out time $\Delta T(\theta_v)$ is given by

$$\Delta T(\theta_v) = 9\Delta t_s(\frac{\gamma_s}{100})^2(\frac{\theta_v}{0.03})^2\frac{S'^2}{S'^2 - 1} \quad (4)$$

Now let us assume at $t = t_0$, the central engine starts to emit another pair of the same slow and rapid sub-jets toward the line of site ($\theta_v = 0$). Then the catch up time (t_c^0) and the break out time (t_b^0) become $t_c^0 = t_0 + t_c$ and $t_b^0 = t_0 + t_b$, respectively, while the relation of the observed time T to t becomes

$$T(\theta_v = 0) = t_0 + \frac{t - t_0}{2\gamma_r^2}. \quad (5)$$

Then

$$T_c(\theta_v = 0) = t_0 + t_r\frac{1}{S^2 - 1}, \quad (6)$$

while the duration from the catch up time to the shock break out time $\Delta T(\theta_v = 0)$ is given by

$$\Delta T(\theta_v = 0) = \Delta t_s\frac{1}{S'^2 - 1}. \quad (7)$$

In this case the observed frequency ν at the detector is related to ν' in the shock frame as

$$\nu = 400\nu'(\frac{\gamma_{sh}}{200}). \quad (8)$$

This means that the observed emission will be in the gamma ray band if ν' is in the soft X-ray band ($\sim \text{keV}$).

Now the condition for the occurrence of the precursor ($T_c(\theta_v = 0) > T_c(\theta_v)$) is rewritten as

$$\frac{t_0}{t_r} > (9(\frac{\gamma_s}{100})^2(\frac{\theta_v}{0.03})^2 - \frac{1}{S^2})\frac{S^2}{S^2 - 1}. \quad (9)$$

Let us consider a specific numerical example;

Case 1

$$\begin{aligned} \gamma_s = 100, \gamma_r = 200, \gamma_{sh} \sim 200, t_r = 6s, \\ \Delta t_s = 6s, \theta_v = 0.03. \end{aligned} \quad (10)$$

Then

$$\begin{aligned} T_c(\theta_v) = 72s, \Delta T(\theta_v) = 72s, \\ T_c(\theta_v = 0) = t_0 + 2s, \Delta T(\theta_v = 0) = 2s. \end{aligned} \quad (11)$$

If the peak energy in the shock frame is $\sim 0.5\text{keV}$, then the peak energy from 72s to 144s is $\sim 5\text{keV}$ while the peak energy from $t_0 + 2s$ to $t_0 + 4s$ is $\sim 200\text{keV}$. Therefore if $t_0 > 70s$ the X-ray emission starts ($t_0 - 70s$) before the gamma ray emission. The duration of the X-ray precursor is $\sim 70s$ while the duration of the gamma ray emission is $\sim 2s$. The duration of the X-ray precursor in this specific example happens to be comparable to those observed by Ginga for GRB900126 (Murakami *et al.* 1991) and by Beppo/SAX for GRB980519 (in't Zand *et al.* 1999) while the duration of the gamma ray is somewhat smaller.

The above model is extremely simple but clearly demonstrates a possible origin of precursors. In Case 1 by choosing t_0 much smaller than 70s we may explain the post-cursor. For example if we choose $t_0 = 20s$ then the gamma ray emission starts at 22s and ends at 24s while the X-ray post-cursor emission starts at 72s.

3. X-ray flux from the precursor and the post-cursor in a sub-jet model

In the previous section, we discussed only the peak energy and the temporal structure of precursors and post-cursors. In this section we estimate the

X-ray flux from the precursor in the sub-jet model. The observed X-ray flux of the precursor and post-cursor of GRB980519 (in't Zand *et al.* 1999) is $\sim 5 \times 10^{-9} \text{erg s}^{-1} \text{cm}^{-2}$ in 2-10 keV band. The average X-ray flux of the precursor of GRB900126 (Murakami *et al.* 1991) is $\sim 2.5 \times 10^{-9} \text{erg s}^{-1} \text{cm}^{-2}$. We argue that these values are compatible with our model of precursors and post-cursors.

For $\theta_v = 0$ and $\gamma\Delta\theta_\gamma \gg 1$, we assume that the observed spectrum of the gamma ray from the sub-jet is given by

$$F_\nu \propto \frac{1}{(1 + (\frac{\nu}{\nu_0})^2)^{\beta_B/2}}, \quad (12)$$

where $\beta_B \sim 1.5$ and $\nu_0 \sim 150 \text{keV}$ are constants. The above spectrum is essentially the same as the Band spectrum (Band *et al.* 1993) with $\alpha = -1$ and $\beta = -\beta_B - 1$, where α and β are the spectrum parameters in the Band spectrum. The peak frequency ν_{peak} , that is, the maximum of νF_ν , is $\sqrt{1/(\beta_B - 1)}\nu_0$.

Garnot, Piran & Sari 1998 as well as Woods & Loeb 1999 derived a general formula to compute the off-axis emission from beamed GRBs. Here we adopt their formulations and notations. Let us use a spherical coordinate system $\mathbf{r} = (r, \theta, \phi)$ where the coordinate are measured in the lab frame; let the $\theta = 0$ axis (z -axis) points to the detector and $\mathbf{r} = 0$ be the central engine. Let also D be the distance to the source and $\alpha = r \sin \theta / D$ be the angle that a given ray makes with the normal to the detector. Then the observed flux is given by

$$F_\nu(T) = \frac{\nu D}{\gamma \beta} \int_0^{2\pi} d\phi \int_0^{\alpha_m} \alpha^2 d\alpha \times \int_{\nu\gamma(1-\beta)}^{\nu\gamma(1+\beta)} \frac{d\nu'}{\nu'^2} \frac{j'_{\nu'}[\Omega', \mathbf{r}, T + \frac{r\mu}{c}]}{\{1 - \mu^2\}^{3/2}}, \quad (13)$$

$$\mu = (1 - \nu'/\gamma\nu)/\beta, \quad (14)$$

$$\gamma = 1/\sqrt{1 - \beta^2}, \quad (15)$$

where α_m , T and $j'_{\nu'}$ are the maximum value of α , the arrival time of a photon at the detector and the rest frame emissivity measured in $\text{erg s}^{-1} \text{cm}^{-3} \text{Hz}^{-1} \text{sr}^{-1}$, respectively. Note here that $'$ means the physical quantity in the rest frame.

In the case of $\theta_v = 0$, we assume that $j'_{\nu'}$ is expressed as

$$j'_{\nu'} \propto H(\Delta\theta_\gamma - \theta) f(\nu'), \quad (16)$$

where $f(\nu')$ and $H(x)$ are a certain function and the Heaviside step function, respectively. Then $F_\nu(T)$ is

given by (Nakamura 1999)

$$F_\nu(T) = 2\pi A \nu \int_{\nu/2\gamma}^{\nu\gamma\Delta\theta_\gamma^2/2} f(\nu') \frac{d\nu'}{\nu'^2}, \quad (17)$$

where A is a factor depending on the density, the gamma factor, the strength of the magnetic fields and the location of the shock as well as the distance to the source. Since we assume that the spectrum has the form of Eq. (12) for $\gamma\Delta\theta_\gamma \gg 1$, $f(\nu')$ is given by

$$f(\nu') = \frac{1 + (\beta_B + 1)(\frac{2\gamma\nu'}{\nu_0})^2}{2\gamma(1 + (\frac{2\gamma\nu'}{\nu_0})^2)^{\beta_B/2+1}}. \quad (18)$$

Neglecting the term proportional to $1/(\gamma\Delta\theta_\gamma)^2$ we have

$$F_\nu(T) = 2\pi A \frac{1}{(1 + (\frac{\nu}{\nu_0})^2)^{\beta_B/2}} \quad (19)$$

Then the peak frequency is in the gamma ray region as $\nu_{peak} \sim \nu_0 \sim 150 \text{keV}$ and the peak flux is given by

$$(\nu F_\nu)_\gamma \sim 2\pi A \nu_0 / 2^{\beta_B/2}. \quad (20)$$

Now let us consider the off axis case ($\theta_v > \Delta\theta_\gamma$). In this case $j'_{\nu'}$ is expressed as

$$j'_{\nu'} \propto H(\cos \phi - (\frac{\cos \Delta\theta_\gamma - \cos \theta_v \cos \theta}{\sin \theta_v \sin \theta})) \times H(\Delta\theta_\gamma - |\theta - \theta_v|) f(\nu'), \quad (21)$$

The flux is given by (Nakamura 1999)

$$F_\nu(T) = 2A \nu \int_{\nu\gamma(1-\beta \cos(\theta_v - \Delta\theta_\gamma))}^{\nu\gamma(1-\beta \cos(\theta_v + \Delta\theta_\gamma))} \phi_v f(\nu') \frac{d\nu'}{\nu'^2}, \quad (22)$$

where $\cos \phi_v = (\cos \Delta\theta_\gamma - \cos \theta \cos \theta_v) / \sin \theta_v \sin \theta$.

Eq.(22) is evaluated as

$$F_\nu(T) = 8A (\frac{\Delta\theta_\gamma}{\theta_v})^2 (\frac{1}{\gamma\theta_v})^2 2\gamma f(\nu\gamma\theta_v^2/2). \quad (23)$$

Then the peak frequency is in the X-ray region as $\nu_{peak} \sim \nu_0/(\gamma\theta_v)^2 \sim 4(\gamma/200)^{-2}(\theta_v/0.03)^{-2} \text{keV}$ and the peak flux is given by

$$(\nu F_\nu)_X \sim 8A (\frac{\Delta\theta_\gamma}{\theta_v})^2 (\frac{1}{\gamma\theta_v})^4 \frac{2 + \beta_B}{2^{\beta_B/2+1}} \quad (24)$$

The ratio of the X-ray flux observed from the off axis of the sub-jet to the gamma ray flux observed along the axis is given by

$$\frac{(\nu F_\nu)_X}{(\nu F_\nu)_\gamma} = 10^{-3} (1 + \beta_B/2) (\frac{\Delta\theta_\gamma}{\theta_v})^2 (\frac{\gamma}{200})^{-4} (\frac{\theta_v}{0.03})^{-4}. \quad (25)$$

For the gamma ray flux of $\sim 5 \times 10^{-6} \text{erg s}^{-1} \text{cm}^{-2}$, we have the X-ray flux of $\sim 5 \times 10^{-9} \text{erg s}^{-1} \text{cm}^{-2}$. This X-ray flux is similar to the observed ones in the precursors of GRB980519 (in't Zand *et al.* 1999) and GRB900126 (Murakami *et al.* 1991). Therefore our model is compatible with the X-ray flux of the precursor and the post-cursor.

4. Discussion

Let us try to interpret the X-ray photon index evolution of GRB980519 (in't Zand *et al.* 1999) stated in Introduction. In our model X-ray precursor and post-cursor come from the off-axis emission from sub-jets so that the spectrum is expressed by Eq. (23) which leads to the photon index $\beta = -\beta_B - 1 \sim -2.5$. While the main gamma ray emission comes from the on-axis emission from the sub-jet so that the spectrum is expressed by Eq. (19). In Eq.(19) the photon index in the X-ray band is -1 . Therefore our model predicts that the photon index evolves from -2.5 to -1 to -2.5 which agree with the observations.

As we discussed in section 2, the duration of the emission from the sub-jet depends on the viewing angle. From Eqs. (4) and (7), we have

$$\frac{\Delta T(\theta_v)}{\Delta T(\theta_v = 0)} = 36 \left(\frac{\gamma_s}{100} \right)^2 \left(\frac{\theta_v}{0.03} \right)^2 \left(\frac{S'}{2} \right)^2. \quad (26)$$

This means that the duration of the lower energy emission ($\theta_v \neq 0$) is longer than that of the higher energy emission ($\theta_v = 0$). If the X-ray precursor or post-cursors occur by chance, our model in this *Letter* predicts that their duration are longer than the duration of the main gamma ray emission. This agrees with observations of precursors and post-cursors (Murakami *et al.* 1991, in't Zand *et al.* 1999).

However for most cases precursors and post-cursors do not occur. Then what does our model predict for usual GRBs? In our model, in general, there exists a sub-jet whose axis makes an angle $\theta_v \neq 0$ with the normal to the detector. This sub-jet mainly emits X-ray delaying from the main gamma ray emission from the sub-jet with $\theta_v \sim 0$. The duration of X-ray emission is longer than the main gamma ray emission so that our model predicts that prompt and delayed X-ray emission with longer duration should associate with GRBs. The emission patterns of GRB960720, 970111, 970228, 970508, 971214, 980329 and 980425 qualitatively agree with this prediction (Frontera *et al.* 1999) while other possible explanations may exist.

What is a new prediction from our model? Suppose that by chance none of the sub-jets point to the line of sight. In this case gamma ray emission is extremely weak and short or absent while the X-ray precursors and/or post-cursors as well as X-ray, optical and radio afterglow exist. If such an event is found, that will be compatible with our sub-jet model discussed in this *Letter*.

I am very grateful Murakami, Kawai and Yoshida for useful comments. This work was supported in part by Grant-in-Aid of Scientific Research of the Ministry of Education, Culture, and Sports, No.11640274 and 09NP0801.

REFERENCES

- Band, D. *et al.* 1993, ApJ, 413, 281
- Castro-Tirado, A.J. *et al.* 1999, Science, 283, 2069
- Dai, Z. G. & Lu, T. 1999, ApJ, 519, L155
- Garnot, J., Piran, T. & Sari, R. 1998, ApJ, 513, 679
- Frontera, F. *et al.* 1999, astro-ph/9911228
- Fruchter, A.S. *et al.* 1999, ApJ, 519, L13
- Galama, T.J. *et al.* 1999, astro-ph/9903021 (nature in press)
- Halpern, J. P., Kemp, J., Piran, T. & Bershadsky, M. A. 1999, ApJ, 517, L105
- Harrison, F. A. *et al.* 1999, ApJ, 523, L121
- Kumar, P. & Piran, T. 1999 astro-ph/9909014
- Kulkarni, S.R. *et al.* 1999, astro-ph/9902272
- Murakami, T. *et al.* 1991, Nature, 350, 592.
- Nakamura, T. 1999, ApJ, 522, L101
- Paczynski, B. 1998, ApJ, 494, L45
- Piran, T. 1998, Phys. Rep. 314, 575 and references therein
- Rhoads, J.E. 1997, ApJ, 478, L1
- Sari, R., Piran, T. & Narayan, R., 1998, ApJ, 497, L17
- Sazonov, S. Y. *et al.* 1998, AAS, 129, 1.

Stanek, K. Z. *et al.* 1999, ApJ,522 , L39

Woods, E. & Loeb, A. 1999, astro-ph/9903377

in't Zand, J. J. *et al.* 1999, ApJ,516 , L57